




RED II System

Rigid External Distraction



Oral and maxillo-facial surgery is our passion! Its further development, together with our customers, is our ambition. Every day we work on developing innovative products and services which meet the highest demands on quality, and which contribute to the wellbeing of the patient.

Rigid External Distraction **RED II** System

Most patients showing midfacial hypoplasia are usually preoperated. Often, a large amount of scar tissue formation is limiting the success of any distraction procedure ending up in compromising results. There the RED II is definitely setting new standards. It is extremely efficient in bringing the bone segments in the desired position and simultaneously to keep them there for bone consolidation. As all important components are external, the important vector planning can be corrected at any time. A wide selection of accessories is at your disposition to match any clinical task.

With the introduction of the RED frame back in 1995 KLS Martin was a pioneer company to offer an external halo frame for the correction of severe maxillary hypoplasia mostly associated with Cleft Lip and Palate (CLP) patients.

The incredible successful treatment outcomes led to a complete redesign of the now called RED II frame back in 2000. Since then, the device has been lighter, but simultaneously stronger and more flexible in its application. Over the years the increasing demand of doctors for patient specific solutions led to a bunch of new products. It is the aim of this leaflet to introduce all these modifications to the public.


Product benefits

- Completely adjustable for any midfacial hypoplasia patient
- Possibility to perform Le Fort I, II, III and monobloc distraction procedures
- Force application only on the affected treatment region
- External distractor – easy definition and correction of all vectors at any time
- Unlimited distraction distances
- Very strong distraction force, excellent retention potential
- Easy and quick assembly in the OR as well as removal in the office or clinical setting
- Ability to treat patients with severe skeletal deficiencies who are not amenable to, or would receive compromised results with conventional orthognathic surgery
- No bone grafting required – no uncalculable recidiva involved

Rigid External Distraction



RED II System

For the usual Le Fort I procedure, 51-580-00-04 is already providing most of the items needed. The listing below shows you what it takes.



RED II Distraction system

Item No.			
51-580-00-04			RED II Distraction system, complete assembly
Consisting of:			
1	51-580-01-04	1	Distraction segment, left
2	51-575-15-04	2	Carbon rods, 120 mm, horizontal
3	51-580-05-04	1	Center part
4	51-575-16-04	1	Carbon rod, 150 mm, vertical
5	51-580-45-04	1	Horizontal cross bar assembly, complete with horizontal cross bar + holder + 2 spindle units
6	51-580-02-04	1	Distraction segment, right
7	51-580-85-07	1	Patient screwdriver
To order separately:			
8	51-575-90-07	1	Adjustment screwdriver, hexagonal
9	51-575-10-09	1 Pack	Fixation screws 45 mm, 10/each
or			
	51-575-12-09	1 Pack	Fixation screws 55 mm, 10/each




What do you need
for which procedure?



Item No.	Description	Unit	Qty (per pack)
1) Must for Le Fort I and Le Fort II procedures			
51-580-00-04	RED II complete, also containing the patient screwdriver	1	1 piece
51-575-90-07	Hexagonal screwdriver (for adjustment and pin fixation	1	1 piece
51-575-10-09	Fixation screws 45 mm for the adult patient	1	10 pcs.
or 51-575-12-09	Fixation screws 55 mm for the pediatric patient	1	10 pcs.
2) Connection to the occlusal level			
Either	Intraoral splint for connecting the RED to the teeth as shown on pages 11-13 in this brochure	1	1 piece
or	Retention plates as presented on pages 14-21 in this brochure		
	General recommendation: 2 pcs. 51-582-50-04 (1.5-mm system)	2	1 piece
3) Additionally for Le Fort III and monobloc procedures			
all items as listed under 1) + 2) and additionally			
51-580-45-04	Second horizontal cross bar	1	1 piece
51-581-02-09	Threaded fixation plate	2	1 piece
51-581-15-09	Threaded fixation pin, 15 mm long (see page 25)	2	1 piece
51-500-90-07	Patient screwdriver straight	1	1 piece
25-665-05-09	Centre Drive® screws 1.5 x 5 mm	1	5 pcs.
to 25-665-07-09	Centre Drive® screws 1.5 x 7 mm (equivalent maxDrive® screws would also be correct)	1	5 pcs.
25-402-99-07	Screwdriver handle	1	1 piece
25-430-98-07	Blade for 1.5-mm Centre Drive® screws	1	1 piece

Standard set see pages 4-5



Adjustment of the RED II frame



Item No	51-580-85-07	51-575-90-07	51-500-90-07	25-402-99-07 Screwdriver handle 25-489-97-07 Blade for 1.5-mm maxDrive® screws 25-430-98-07 Blade for 1.5-mm Centre Drive® screws
Application	Patient screwdriver hexagonal	Adjustment screwdriver hexagonal	Patient screwdriver straight	Screwdriver 1.5 mm Micro
Description	Activates the distraction movement	For all intraoperative adjustments of the RED II-frame 2 working ends	For insertion of the threaded insertion pin in LeFort III and monobloc procedures	For fixation of the threaded fixation plate

Patient-specific **RED II** with temporal fixation plates



The standard RED II frame is mostly efficient in very young children. The fixation of the conventional titanium pins however remains a challenging task.

Therefore, as an alternative to our standard RED II a patient-specific design according to the ideas of Prof. J. Obwegeser (Limmatklinik Zurich, Switzerland) is available.

The RED II according to Prof. J. Obwegeser comes with the standard head frame being fixed to temporal fixation plates that are individually manufactured to the anatomics of each patient.


This custom-made device (no CE marking) has to be requested for every single patient via our IPS Gate(R) upload and communication portal.

For further information visit the KLS Martin website.



Benefits of the temporal fixation plates and the IPS® Planning Service:


- Stronger anchoring to the skull bone
→ Reduced risk of the RED II distractor becoming detached in the event of the patient falling
- Application in patients with thin-walled skull bone or poor bone quality
- Application in patients with openings in the cranial bone in the region of the temporal bone
- Hook which passes around the zygomatic bone for easier positioning
- If necessary, planning & simulation of the post-operative situation after distraction (IPS® Planning Service)
- If necessary, modified fixation and retention plates in the midface available as IPS® products
- With the help of 3D visualisation a heatmap of the bone thickness in the region of the temporal fixation plates allows for more precise planning



Connection on the Le Fort I-Level Via Retention plates or via Intraoral Splint

Intraoral Splint

= Tooth-borne attachment to the maxilla
(see pages 11-13)




Retention Plates

= Bone-borne attachment to the maxilla
(see pages 14-21)






The completed splint is cemented in the clinical setting and at the time of surgery circumdental wires are passed through most of the maxillary teeth to increase stability.



Reinforced external traction hook in a preoperated patient. A piece of wire is soldered diagonally to decrease the cantilever effect at the free end of the hooks.



Completed intraoral appliance – the outer bow has been bent to form the traction hooks. Note small soldered hooks to be used during the facial mask retention phase after distraction.

The Intraoral Splint

In order to apply traction to the maxilla through dentition, a rigid intraoral splint is often the most adequate option.

Orthodontic bands with 0.045 to 0.050 inch head-gear tubes are fitted either on the second primary molars (children under 6 years) or the first permanent molars and an alginate or compound impression is taken of the maxillary arch.

The bands are transferred and the impression is poured with dental stone. The splint is made on the working model. If the patient does not have orthodontic brackets, the labial and palatal wires are bent in close contact with most of the maxillary teeth. If the patient has orthodontic brackets, the labial wire has to be bent outward and gingivally to clear the existing appliances. If needed, a trans-palatal bar can be added to increase rigidity. Connecting wires between the labial and palatal arches through the embrasures between the lateral and canine teeth bilaterally or in any other area where the wire can be passed without interfering with the occlusion may also be incorporated.

The device is inserted just prior to OR at the time of surgery. It is preferable to do maxillary arch expansion procedures before or after distraction to avoid moving the maxillary bone simultaneously in several directions where vector control can become more difficult. If the clinician desires to expand simultaneously with anterior distraction, an expansion screw can be incorporated into the splint, which has to be split into two segments. The stability of the device may then be compromised.

The intraoral splint is not a KLS Martin standard product. It will be manufactured by the hospital's orthodontic team. Individual differences on patient's dentation may demand a different orthodontic splint.



Fig. 1:
9 year 10 month old boy with a repaired left unilateral cleft lip and palate presented with severe maxillary hypoplasia.



Fig. 2:
The preoperative facial photographs demonstrate the mid-face deficiency with a concave facial profile and retrusive upper lip.



Fig. 3:
After maxillary distraction the facial profile and balance were restored to normal proportions.



Fig. 4:
Note the improved prominence at the malar level and the improved relationship between the upper and lower lips. Nasal form was also improved as a result of the maxillary advancement through distraction osteogenesis.




Fig. 5:
Intraorally there were marked anterior and bilateral posterior crossbites.




Fig. 6:
The postoperative intraoral photographs demonstrate complete correction of the anterior crossbite. One year after distraction the patient has not shown signs of relapse.






Fig. 7:
The patient underwent a high two piece Le Fort I osteotomy with pterygomaxillary and septal disjunctions. No bone grafting or rigid internal fixation hardware was utilized. There was no repositioning of the maxilla at the time of the surgery.

The RED device was placed immediately after the osteotomy and the patient was discharged the morning after surgery. Distraction was initiated on postoperative day 5 at the rate of 1 mm per day.

The total maxillary advancement was 10 mm. Three weeks of rigid retention were utilized.



Connection on the Le Fort I-Level Via Retention plates



Impact of the point of anchorage on the rotational movements of the midface during distraction


The chart shows the impact of various fixation points on the maxilla. An anchorage on the tooth level will usually lead to a posterior rotation, which is often not desired. An anchorage point higher up, in the center of resistance or above would be better, because this would lead to none or to an anterior rotation.

The advantages are


- Solid bone fixation where high distraction forces are involved
- Minor risk of periodontal harm or teeth extractions
- Ready-made – no need for the orthodontist to customize the wire bar
- Easier dental hygiene compared to orthodontic band fixation
- Accurate distraction vector setting, no unwanted rotational movements
- Easy fixation and removal
(as in standard osteosynthesis plates)

Contraindications


- Cases of inadequate bone volume to fix the osteosynthesis plate.
The general rules and guidelines of Distraction Osteogenesis have to be followed.



Dental anchorage may cause counter-clockwise rotation of the maxilla.




Anterior position of the retention plate will lead to a spring-like action. → Posterior rotation and vertical midfacial elongation.



A posterior placement of the retention plate will counteract the posterior vertical elongation. The surgeon will need long quadrangular rods and a posteriorly-placed rider.


Further indications for retention plates are:

- Distraction to be performed on edentulous patients or patients with severe periodontal disease or the existing risk of periodontal damage
- Especially Cleft Lip and Palate (CLP) patients can often only offer a limited dentition for dental anchorage
- If the maxilla is not only moved horizontally, but also vertically in a downwards direction there is a danger of pulling the wire fixation off the teeth
- Left and right maxillary segments can be manipulated independently which is a major benefit especially in Cleft Lip and Palate (CLP) patients
- Even multipiece distractions (e.g. 3 segments) can be performed
- Simultaneous rapid maxillary expansion is possible (f.e. transversal distraction can be performed during procedure)
- Retention plates are a prerequisite for sutural midfacial distraction



Retention plates and retention plate connectors

Leipzig Retention Plate*



Retention plates

Item Numbers:

51-582-50-04	Set 1.5 mm complete (1 each)
51-582-55-04	Set 1.8 mm complete (1 each)

To be fixed with 1.5 mm screws

Set includes:

1 bone plate, 11 holes
1 rider incl. screws for rod fixation
1 square rod either 1.8 or 1.5-mm thick
1 fixation eyelet


The entire set is designed for single use only !

To be modified using a 1.5-mm maxDrive® screwdriver



* Developed in cooperation with PD Dr. Dr. Thomas Hierl / Prof. Dr. Dr. Alexander Hemprich, Leipzig, Germany

New items:



51-582-51-09



Solidly connected retention plate:

Between fixation plate and quadrangular rod.

Advantages:

- No connection elements needed
- No risk of losing connection elements
- No risk of harming the patient with exposed metallic elements


Disadvantage:

- No lateral attachment is possible.

To be modified using a 1.5-mm maxDrive® screwdriver
Unit: 1 piece each



51-580-13-09



Retention plate connector

Allows a direct attachment of 1.5- and 1.8-mm retention plate eyelets to the quadrangular rods activation spindles of the RED-frame (2 pcs. each).

To be modified using a 1.5-mm maxDrive® screwdriver
Unit: 1 piece each





Fig. 1:
63-year-old patient suffering from CLP. Only one maxillary molar is left. Referral due to insufficiency to facilitate prosthodontic therapy.




Fig. 2:
Preoperative CT reconstruction. Severe maxillary retrusion and atrophy.



Fig. 3:
Frontal view



Fig. 4:
Preoperative lateral cephalogram. Marked midfacial retrusion, no bone stock for implant insertion or prosthodontic therapy.



Fig. 5:
Lateral cephalogram after distractor removal. As no dental occlusion will stabilize the new midfacial position, miniplates are temporarily inserted. Simultaneously a bilateral sinus lift procedure and bone augmentation in the cleft area was performed. The bent miniplates represent the amount of forward maxillary displacement. Dental implants will be inserted 3 months later.



Fig. 6:
Situation before removal of the RED. See the improvement in midfacial prominence and the uprightening of the nose.




Fig. 7:
Preoperative intraoral situation



Fig. 8:
Situation 3 years after distraction, augmentation and implant insertion. Magnetic abutments are used for prosthesis fixation.



Fig. 9:
Situation 3 years after distraction. Marked esthetic improvement, good facial balance.



Fig. 10:
Lateral cephalogram 3 years after distraction, augmentation and implant insertion.



Fig. 1:
19-year-old man suffering from unilateral Cleft Lip and Palate (CLP). Note the maxillary retrusion and midfacial hypoplasia leading to collapsed and inwardly rotated maxillary segments.



Fig. 2:
Preoperative dental situation



Fig. 3:
Facial profile view, significant malar deficiency.



Fig. 4:
Post-distraction situation. See the alignment of both maxillary segments using Leipzig retention plates. To correct malar asymmetry, the osteotomy line has been extended on the smaller maxillary segment. Bone grafting in the cleft area and paranasal region was performed during distractor removal.



Fig. 5:
Occlusion 4 years after distraction osteogenesis shows stable results. In the meantime, a dental implant has been inserted in the cleft region.







Fig. 6:
Facial profile 4 years after two-piece segmental distraction. See improved facial balance.



Spare parts and variations of the **RED II**



Item No.			
10	51-580-40-07	1	Fixation ring
11	51-580-35-04	1	Holder for horizontal crossbar
12	51-580-97-04	1	Head cap screw
13	51-575-95-04	1	Hexagonal nut
14	51-575-98-07	1	Hexagonal nut



Expansion of the **RED II** additional components



51-580-45-04

Horizontal crossbars:

The redesigned horizontal crossbars and their new spindle units are designed to allow 3D-steering of the distraction movement.

Furthermore, an expansion of the maxilla is now possible as well. Loosen the screw, select new position and lock.

unit: 1 piece



To be modified using 51-575-90-07

Remarks:

For Le Fort II, Le Fort III and Monobloc procedures a second horizontal cross bar is recommended.

1 distraction unit will always come with the basic RED frame configuration, e.g. 51-580-00-04.

51-580-45-04: Horizontal cross bar in **purple color**




51-580-26-04

In order to possibly update existing RED II distraction devices with the new spindle, one can order the spindle as a spare part.

unit: 1 piece

To be modified using 51-575-90-07





Expansion of the **RED II** additional components

Fixation screws



51-575-10-09

Fixation screw 45 mm

Unit: 10 pieces each



51-575-12-09

Fixation screw 55 mm

Unit: 10 pieces each

The longer fixation pin, usually applied for children



51-575-14-09

Trial fixation pin, 41 mm

Unit: 1 piece each

To be used for intraoperative setting of the RED II.

Blunt tips – not for permanent fixation !

To be modified using 51-575-90-07 

Locking nuts and stops



51-575-94-09



51-575-99-09




The **locking nut** 51-575-94-09 is designed to prevent loosening and over-tightening of the fixation pin.


Unit: 1 piece each

The **positive stop** 51-575-99-09 securely limits the skull entry of the RED fixation pin.

Unit: 1 piece each



51-580-08-04



Halo extender

Allows pin fixation on the posterior part of the skull and an extension of the RED-frame. Symmetrical construction – to be used on the right or left side of the patient.

Unit: 1 piece each



51-583-01-04

Rounded fixation element left

Enables the placement of fixation pins on various levels



51-583-02-04


Rounded fixation element right

Enables the placement of fixation pins on various levels



51-583-00-04

RED II with rounded fixation element complete,
according to the specifications on page 4-5



Expansion of the **RED II** additional components

Central fixation pins and fixation plates



Micro screws usually
1.5 x 5 mm to 1.5 x 7 mm



51-581-15-09

51-581-15-09

Threaded central fixation pin, 2.0 x 15 mm:

Unit: 1 piece each

To be inserted using 51-500-90-07



51-581-21-09

51-581-21-09

Threaded central fixation pin, 2.0 x 21 mm:

Unit: 1 piece each

To be inserted using 51-500-90-07



51-581-30-09



51-581-30-09

Threaded central fixation pin, 2.0 x 30 mm:

Unit: 1 piece each


To be inserted using 51-500-90-07






51-581-08-09
Habal type 8-mm pin
 (5 mm threaded)
 Direct anchorage on the affected bone
 Unit: 1 piece each

To be inserted using 51-500-90-07 



51-581-10-09
Habal type 10.5-mm pin
 (7.5 mm threaded)
 Direct anchorage on the affected bone
 To be applied with 51-500-90-07
 Unit: 1 piece each

To be inserted using 51-500-90-07 



51-581-02-09
Straight threaded fixation plate:
 For Le Fort III and Monobloc procedures, a second fixation base allows a better control of the distraction vector and the bony structures involved.
 Unit: 1 piece each

To be fixed with 1.5 mm screws



51-581-03-09
Threaded fixation plate*
 is an alternative to the straight threaded fixation plate 51-581-02-09.
 Unit: 1 piece each




51-581-06-09
Threaded fixation plate*
 (0.5 mm threaded)
 is an alternative to the straight threaded fixation plate 51-581-02-09.
 Especially suitable in round, suborbital bone regions
 Unit: 1 piece each

* All to be applied with 1.5-mm micro screws
 (usually 5 to 7 mm long) on the lateral aspect.


Sutural Midface Distraction

Sutural midfacial distraction (SMD) utilizes the high forces which can be applied with the RED device to a growing organism. Without the need for osteotomies, complex changes of the midfacial architecture may be achieved in short time. It is of paramount importance to check bone thickness of the calvarium prior to SMD to avoid skull punctures or even skull fractures.


Furthermore dental splints must not be used as dental extrusion will result. As SMD is a new procedure, thorough treatment planning and control of the patient during the procedure is mandatory. SMD may not be performed in adult patients.



Retention plates fixed to the midface. Note the bending of the plate to utilize the bone stock of the zygomatic buttress. As anatomy is highly variable, retention plates with moveable riders are suggested.



At least 3 screws anterior to the rider and as many as possible posteriorly should be used. 1.5-mm Drill-Free screws have been inserted. No osteotomy was performed. Standard distraction activation of 1 mm/day is used.




Same patient (6 ys.; syndromal midfacial retrusion) before and after SMD. Midfacial advancement, opening of all sutures (e.g. zygomatic arch), rotation of the midface and rotation of the nasal bones is visible. Due to protraction forces, the maxillary arch will change shape, too.

References

- Hierl, Th.; Klöppel, R.; Hemprich, A.:
Midfacial distraction osteogenesis without major osteotomies – a report on first clinical application
Plast Reconstr Surg 108 (2001), 1667-1672
- Hierl, Th.; Hemprich, A.:
A novel modular retention system for midfacial distraction osteogenesis.
Br J Oral Maxillofac Surg 38 (2000), 623-626
- Hierl, Th.:
Lengthening the maxilla by distraction osteogenesis.
In: Bell, W; Guerrero, C.: Orthognathic surgery vs distraction osteogenesis. Quintessence Int. 2007.

Storage Recommendation



Storage Recommendation

Qty	Item No	Specification
1x	55-804-50-01	Mesh Tray 477 x 251 x 64 mm
1x	55-805-52-01	Lid
1x	55-234-13-04	Marsafe Container 553 x 272 x 133 mm
1x	55-891-40-01	Small-parts basket, fine mesh 80 x 80 x 40 mm
1x	55-806-11-04	3x Instrument holder, Ø 15 mm high
1x	55-806-12-04	3x Instrument holder, Ø 20 mm high
1x	55-806-20-04	3x Fixation element universal H = 40 mm
1x	55-806-10-04	6x Instrument holder Ø 8-10mm high
2x	55-806-50-04	Separator 123 x 9 x 22mm, with clips
7x	55-806-25-04	6x Studded strip, 22 mm

Publications and Literature



Source: Dr. Camilo Roldán, Hamburg

German

P. Kessler, F. Kloss, U. Hirschfelder, F. W. Neukam, J. Wiltfang
Osteodistraktion im Mittelgesicht, Indikation, Technik und
erste Langzeitergebnisse
DFZ 2/2004, S. 1-6

Hierl T., Primm T., Klöppel R., Hemprich A.
Therapie ausgeprägter Mittelgesichtsrücklagen mit Hilfe der
Distraktionsosteogenese
Mund Kiefer GesichtsChir 2003, 1-2003, S. 7 ff

Hierl T., Primm T., Klöppel R., Hemprich A.
Distraktionsosteogenese im Mittelgesichtsbereich.
Grundlagen und klinische Anwendung
Quintessenz 51, 3, S. 247-256, 2000

Hierl T., Primm T., Klöppel R., Hemprich A.
Einsatz der Kallusdistraktion bei ausgeprägter Mittelgesichts-
hypoplasie
Dtsch. Zahnärztliche Zeitung Z 55 (2000), S. 359-362

English

Ahn J-G, Figueroa AA, Braun S, Polley JW:
Biomechanical considerations in distraction of the osteotomized
dentomaxillary complex
Am J Dentofac Orthop 116: 264, 1999

Cheung L.K., Chua H. D.
Maxillary Distraction for Patients with Cleft Lip and Palate (CLP)
In Bell W., Guerrero C. Distraction Osteogenesis of the Midface,
BC Decker 2007, p. 529-542

van Eggermont B., Jansen J., Bierman M.W.J.
Patient satisfaction related to rigid external distraction
osteogenesis,
Int. J. Oral Maxillofac. Surg. 2007; 36; p. 896 - 899

Ghali, G.E., Sinn D.P.
Gradual Repositioning of the Midface at the Sub-cranial Le Fort III
Level by Distraction Osteogenesis
In Bell W., Guerrero C. Distraction Osteogenesis of the Midface,
BC Decker 2007, p. 285-291

A.A. Figueroa, J. W. Polley
Management of severe cleft maxillary deficiency with distraction
osteogenesis: Procedure and results
American Journal of Orthodontics, Vol 5, No.1, March 1999,
p. 46-51

A. A. Figueroa, J. W. Polley, E. Ko
Distraction Osteogenesis of Severe Cleft Maxillary Deficiency
with the RED Technique
In: M. L. Samchukov, J.B. Cope, A.B. Cheraskin: Craniofacial
Distraction Osteogenesis, 2001, p. 485 - 494

Figueroa AA, Polley JW.
Orthodontic procedure for maxillary distraction.
In International Congress on Cranial and Facial Bone Distraction
Processes.

Figueroa, AA, Polley, JW.
Management of severe cleft maxillary deficiency with distraction
Osteogenesis: Procedure and Results.
Amer. J. Orthod. Dentofacial Orthop., 1999; 115-1-12.

Publications and Literature

Figueroa, AA, Polley, JW, Ko, EW-C.
Maxillary distraction for the management of cleft maxillary hypoplasia with a rigid external distraction system.
Seminars in Orthodontics, 1999; 5: 46-51.

Hochban W, Ganss C, Austermann KH
Long-term results after maxillary advancements in patients with clefts
Cleft Palate Craniofac J 30: p. 237, 1993

Ko, EW, Figueroa AA, Guyette, TW, Polley JW, Law, WR.
Velopharyngeal changes after maxillary advancement in cleft patients with distraction Osteogenesis using a rigid external distraction device: A 1-year cephalometric follow-up.
Jour Craniofac Surg, 1999; 10:312-320.

Figueroa A.A. , Polley J.W.
External vs. Internal Distraction Osteogenesis for the Management of severe maxillary hypoplasia: External distraction
J. Oral Maxillofac. Surg. 2008; 66; p. 2598 – 2604

Figueroa A.A. Polley J.W.
Management of severe cleft maxillary deficiency with distraction osteogenesis; procedure and results
Am J Orthod Dentoc Orthop 1999, 115: 1-12

Hierl, Th.; Hemprich, A.
Callus distraction of the midface in severely atrophied maxilla – a case report
Cleft Palate Craniofac. J 36 (1999), p. 457-461

Hierl, Th. Hemprich, A.
A novel modular retention system for midfacial distraction osteogenesis
Br J Oral Maxillofac Surg. (2000) 38, p. 623-626

Ko EW, Figueroa AA, Polley JW
Soft tissue profile changes after maxillary distraction
J Oral Maxillofac Surg 58: 959, 2000

Krimmel M, Cornelius CP, Roser M, Bacher M, Reinert S.
External distraction of the maxilla in patients with craniofacial dysplasia.
J Craniofac Surg (2001) 12: p. 458–463

Mavili M.E.; Vargel I.; Tunçbilek G.
Stoppers in RED II distraction device: is it possible to prevent pin migration?
The Journal of craniofacial surgery 2004; 15(3):p 377-383

Nørholt S. E., Bjerregaard J., Moskilde L
Maxillary Distraction Osteogenesis in a patient with Pycnodysostosis – A case report
Amer. Assoc. of Oral and Maxillofac. Surgeons; 2004; 62; p. 1037-1040

Nout E., Wolvius B., van Andrichem L.N.A., Ongkosuwito E.M., van der Wal K.G.H.
Complications in maxillary reconstruction using the RED II device – A retrospective analysis of 21 cases
Int. J. Oral Maxfac. Surg. 2006; 35; p. 897-902

B. L. Padwa
Combined Push-Pull Midface Distraction Osteogenesis
In Bell W., Guerrero C. Distraction Osteogenesis of the Midface 2007, BC Decker 2007, p. 293-298

Polley, J.W., Figueroa, AA.
Management of Severe Maxillary Deficiency in Childhood and Adolescence through Distraction Osteogenesis with an External, Adjustable, Rigid Distraction Device
The Journal of Craniofacial Surgery, 8, (3) 181-185, May 1997.

Polley, J.W., Figueroa, AA.
The Management of Cleft Maxillary Hypoplasia with (RED) Rigid External Distraction. Proceedings of the International Congress on Distraction Osteogenesis of the Facial and Cranial Bones Paris, France June 19 – 21, 1997. 255-260.

Polley, J.W., Figueroa, AA., Hong, KF., Huang, CS
Distraction Osteogenesis in the Treatment of Cleft Maxillary Deformities.
Plastic Surgical Forum XX 127-131, 1997.



Polley, J.W., Figueroa, AA.
Midface Osteodistraction-Commentary on Midface Advancement by Bone Distraction and Distraction Osteogenesis and its Application to the Midface and Bony Orbit in the Craniosynostosis Syndromes.
The Journal of Craniofacial Surgery. 9, (2) 119-122, March 1998.

Polley, J.W., Figueroa, AA.
Rigid External Distraction (RED): It's application in cleft maxillary deformities.
The Journal of Plastic and Reconstructive Surgery, 102 (5). 1360-1372. October 1998.

Polley, J.W., Ko, E.W., Figueroa, A.A., Guyette, T.W., Law, W.R.
Velopharyngeal Changes After Maxillary Advancement in Cleft Patients with Distraction Osteogenesis Using a Rigid External Distraction Device: A 1-Year Cephalometric Follow-up.
The Journal of Craniofacial Surgery, 1999; 10:4:312-320.

Polley, J.W.
Commentary on Maxillary Distraction in Cleft Lip and Palate (CLP) patients: A Review of Six Cases.
The Journal of Craniofacial Surgery, 1999; 10:4:329.

Polley, J.W., Figueroa, A.A.
Maxillary Distraction Osteogenesis with Rigid External Distraction.
Atlas of the Oral and Maxillofacial Surgery Clinics of North America, 1999; Volume 7:1.

Polley J. W., Figueroa AA, Charbel FT, et al
Monobloc craniomaxillofacial distraction osteogenesis in a newborn with severe craniofacial synostosis; a preliminary report,
J. Craniofac Surg 6: 421, 1995

Posnick JC, Dagys AP:
Skeletal stability and relapse patterns after Le Fort I maxillary osteotomy fixed with miniplates: The unilateral Cleft Lip and Palate (CLP) deformity
Plast Reconstr Surg 94: P.924 ff, 1994

S. Reinert, M. Kimmel, C.-P. Cornelius, M. Roser, M. Bacher
Rigid External Distraction of the Maxilla: Technique and Clinical Cases
In: M. L. Samchukov, J.B. Cope, A.B. Cheraskin:
Craniofacial Distraction Osteogenesis, 2001, p. 501 - 494

Suzuki E.Y., Buranastidporn B., Ishii M.
New fixation for Maxillary Osteogenesis using locking attachments
Amer. Assoc. of Oral and Maxillofac. Surg
J. Oral Maxillofac Surg; 64; 2006; p. 1553 -1560

H. C. Schwartz, J. Beumer III
Three Dimensional Midface Distraction
In: M. L. Samchukov, J.B. Cope, A.B. Cheraskin:
Craniofacial Distraction Osteogenesis, 2001, p. 506 - 511

Ueki K., Marukawa K., Nakagawa K., Yamamoto E.
Multidirectional distraction osteogenesis for Crouzon syndrome: A technical note
Int. J. Oral Maxillofac. Surg. 2005; 34, p. 82-84

Varol A., Sencimen M., Sabuncuoglu F., Ölmez H., Basa S.
Maxillary distraction osteogenesis for a patient with pycnodysostosis by rigid external distraction II midface distraction system
Int. J. of Oral and Maxillofacial Surgery, Volume 38, Issue 5, 457-457

Witherow H, Dunaway D, Ponniah A, Hayward R
Monobloc distraction in an infant, using the rigid external distractor: Problems and solutions-A case report.
Journal of cranio-maxillofacial surgery: 36(1):15-20, 2008 Jan


Yamuchi K., Mitsugi M., Takahashi T.
Maxillary Distraction Osteogenesis using Le Fort I osteotomy without intraoperative down-fracture
Int. J. Oral Maxillofac. Surg. 2006; 35; p.493 – 497

CMF Surgery

SonicWeld Rx®

Resorbable implants for use in craniomaxillofacial osteosynthesis

- Resorb x®
- Resorb xG




It is the face that makes humans unique and unmistakable –
“There is nothing that more closely reflects the life of an individual than the human face*.”

Our objective is to simplify craniofacial surgery with specially designed implant systems that ensure optimum satisfaction for both surgeon and patient. Together with renowned users we translate new ideas into innovative products and are constantly enhancing them.

Our range of products includes everything necessary for modern craniofacial surgery. We not only set standards but we also go beyond to take advantage of modern technology in the development of solutions customized for the individual patient.

KLS Martin – your competent and reliable partner for both everyday challenges and special challenges.



* © Kurt Haberstich (*1948)

Distractors

Devices for use in correction of malformations

- Cranial distraction
- Midface distraction
- Mandibular distraction



Individual Patient Solutions

Patient-specific solutions for use in craniomaxillofacial surgery

- IPS Implants®
- IPS CaseDesigner®
- IPS Gate®



LevelOne Fixation

Titanium implants and instruments for use in craniomaxillofacial osteosynthesis

- Traumatology
- Reconstruction
- Orthognatic surgery



App for CMF products

All important information about the CMF portfolio at one glance.



KLS Martin Group

KLS Martin Australia Pty Ltd.

Sydney · Australia
Tel. +61 2 9439 5316
australia@klsmartin.com

KLS Martin do Brasil Ltda.

São Paulo · Brazil
Tel. +55 11 3554 2299
brazil@klsmartin.com

**KLS Martin Medical (Shanghai)
International Trading Co. Ltd.**

Shanghai · China
Tel. +86 21 5820 6251
china@klsmartin.com

KLS Martin India Pvt Ltd.

Chennai · India
Tel. +91 44 66 442 300
india@klsmartin.com

Martin Italia S.r.l.

Milan · Italy
Tel. +39 039 605 67 31
italia@klsmartin.com

Nippon Martin K.K.

Tokyo · Japan
Tel. +81 3 3814 1431
nippon@klsmartin.com

KLS Martin SE Asia Sdn. Bhd.

Penang · Malaysia
Tel. +604 505 7838
malaysia@klsmartin.com

KLS Martin de México S.A. de C.V.

Mexico City · Mexico
mexico@klsmartin.com

Martin Nederland/Marned B.V.

Huizen · The Netherlands
Tel. +31 35 523 45 38
nederland@klsmartin.com

Gebrüder Martin GmbH & Co. KG

Moscow · Russia
Tel. +7 499 792-76-19
russia@klsmartin.com

KLS Martin Taiwan Ltd.

Taipei 106 · Taiwan
Tel. +886 2 2325 3169
taiwan@klsmartin.com

Gebrüder Martin GmbH & Co. KG

Dubai · United Arab Emirates
Tel. +971 4 454 16 55
middleeast@klsmartin.com

KLS Martin UK Ltd.

London · United Kingdom
Tel. +44 1189 000 570
uk@klsmartin.com

KLS Martin LP

Jacksonville · Florida, USA
Tel. +1 904 641 77 46
usa@klsmartin.com

Gebrüder Martin GmbH & Co. KG**A company of the KLS Martin Group**

KLS Martin Platz 1 · 78532 Tuttlingen · Germany
P.O. Box 60 · 78501 Tuttlingen · Germany
Tel. +49 7461 706-0 · Fax +49 7461 706-193
info@klsmartin.com · www.klsmartin.com